

RESEARCH ARTICLE

Higher muscle fiber conduction velocity and early rate of torque development in chronically strength-trained individuals

 A. Del Vecchio,^{1,4}  F. Negro,² D. Falla,³ I. Bazzucchi,¹  D. Farina,⁴ and F. Felici¹

¹Department of Movement, Human and Health Sciences, University of Rome “Foro Italico,” Rome, Italy; ²Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy; ³School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, United Kingdom; and ⁴Neuromechanics and Rehabilitation Technology Group, Department of Bioengineering, Imperial College London, London, United Kingdom

Submitted 9 January 2018; accepted in final form 18 July 2018

Del Vecchio A, Negro F, Falla D, Bazzucchi I, Farina D, Felici F. Higher muscle fiber conduction velocity and early rate of torque development in chronically strength-trained individuals. *J Appl Physiol* 125: 1218–1226, 2018. First published July 19, 2018; doi: 10.1152/jappphysiol.00025.2018.—Strength-trained individuals (ST) develop greater levels of force compared with untrained subjects. These differences are partly of neural origin and can be explained by training-induced changes in the neural drive to the muscles. In the present study we hypothesize a greater rate of torque development (RTD) and faster recruitment of motor units with greater muscle fiber conduction velocity (MFCV) in ST compared with a control cohort. MFCV was assessed during maximal voluntary isometric explosive contractions of the elbow flexors in eight ST and eight control individuals. MFCV was estimated from high-density surface electromyogram recordings (128 electrodes) in intervals of 50 ms starting from the onset of the electromyogram. RTD and MFCV were computed and normalized to their maximal voluntary torque (MVT) values. The explosive torque of the ST was greater than in the control group in all time intervals analyzed ($P < 0.001$). The absolute MFCV values were also greater for the ST than for controls at all time intervals ($P < 0.001$). ST also achieved greater normalized RTD in the first 50 ms of contraction [887.6 (152) vs. 568.5 (148.66)%MVT/s, mean (SD), $P < 0.001$] and normalized MFCV before the rise in force compared with controls. We have shown for the first time that ST can recruit motor units with greater MFCV in a shorter amount of time compared with untrained subjects during maximal voluntary isometric explosive contractions.

NEW & NOTEWORTHY Strength-trained individuals show neuromuscular adaptations. These adaptations have been partly related to changes in the neural drive to the muscles. Here, we show for the first time that during the initial phase of a maximal isometric explosive contraction, strength-trained individuals achieve higher levels of force and recruit motor units with greater conduction velocities.

explosive force contractions; motor unit conduction velocity; motor unit recruitment; neuromuscular assessment; size principle

INTRODUCTION

The human neuromuscular system has the ability to develop high forces in short time intervals (1). It takes ~150 ms to reach high levels of force [$>70\%$ maximal voluntary force (MVF)] during a single-joint maximal voluntary isometric explosive

contraction (7, 20, 34, 51). Explosive force is commonly measured during specific time intervals from the contraction onset or characterized by the slope of the joint torque-time curve [i.e., the rate of torque development (RTD)] during the first 200 ms of force generation (1). Over the past decades there has been an increasing interest in the determinants of explosive force especially in relation to the implications for enhancing athletic performance and for the prevention of falls and injuries (1, 6, 13, 34, 39, 45, 51). Moreover, the RTD has been identified as an important parameter to detect changes in neuromuscular function in addition to the MVF (39).

At the neuromuscular level, RTD is associated with the neural drive to muscles during the very early phase (first ~50 ms) of the contraction (14, 16, 18, 34, 54). The neural drive to muscles is the ensemble of motor neuron action potentials that reach the muscle per unit time (28). During rapid “ballistic” contractions, the size principle of motor unit recruitment is maintained (15); however, the motor unit recruitment thresholds decrease, and discharge rates increase compared with slow-force contractions (16).

There is a strong association between the neural activity during the early phase of muscular contraction (0–50 ms) and explosive force (1, 13, 14, 34, 51). The greater RTD in the first 50 ms of the contraction observed in power athletes is partly associated with a greater electromyogram (EMG) amplitude with respect to controls (51). Moreover, 12 wk of isometric explosive training significantly increase the neural responses during explosive contractions compared with isometric strength training (6). However, the specific neural adaptations in chronically strength/power-trained individuals are largely unknown. The present evidence from cross-sectional studies that compared trained individuals with a control cohort (33, 51) relies on EMG features that are separated from the neural drive to the muscle (10, 11, 23).

At the motor unit level, short-term ballistic training increased motor unit discharge rate of the first four detected motor unit action potentials during isometric ballistic contractions (55). Similarly, 6 wk of strength training increased the motor unit discharge rate during contractions at forces below 30% of maximum (56). It can be hypothesized that chronic strength training involving explosive tasks may also result in a faster recruitment of larger, fast-twitch motor units. A faster motor unit recruitment would increase the explosive force of a muscle because the size of the motor unit (i.e., the recruitment threshold; 37) is associated with the motor unit mechanical

Address for reprint requests and other correspondence: F. Felici, Dept. of Movement, Human and Health Sciences, Univ. of Rome “Foro Italico,” Piazzale de Bosis 6, 00135 Rome, Italy (e-mail: francesco.felici@uniroma4.it).

properties (peak force and force risetime; 36, 53). However, direct or indirect data on motor unit recruitment strategies in chronically resistance-trained individuals are lacking (19). Recently, Methenitis et al. reported significant associations between muscle fiber conduction velocity (MFCV) and rate of force development in strength-trained (ST) individuals (43). However, the estimates of MFCV were obtained during electrical stimulation of individual muscle fibers (43); thus, the time course of MFCV (when the muscle fibers are activated by the central nervous system) is unknown.

There are methodological limitations in the identification of individual motor unit activities in time intervals of 20–50 ms during explosive force contractions. However, it is possible to indirectly assess the properties of active motor units by measuring their average MFCV. MFCV is a size principle parameter (4) since muscle fibers of high-threshold motor units have greater diameters than those of lower-threshold motor units (4, 11, 41). There is a biophysical relation between MFCV and fiber diameter (44, 46) that has been demonstrated at the individual muscle fiber level (35). Moreover, we have recently shown that the increase in the average MFCV during increasing-force contractions is strongly associated with the increase in single-motor unit conduction velocities when related to the progressive recruitment of motor units (10). It has also been shown that MFCV can be reliably estimated from EMG signals in intervals as short as ~25 ms (9, 27, 29, 41).

Therefore, estimates of MFCV may provide an indirect analysis of motor unit recruitment (4, 10, 11). Moreover, the time course of MFCV during maximal voluntary isometric explosive contractions is unknown. In this study, we measure for the first time MFCV during explosive voluntary force contractions in chronically ST individuals compared with untrained subjects. We hypothesized a greater early RTD in ST individuals that would be accompanied with a higher MFCV during isometric explosive force contractions.

MATERIALS AND METHODS

Participants

Sixteen healthy, nonsmoking young men volunteered for this study, which was approved by the University of Rome Ethical Advisory Committee and conducted according to the Declaration of Helsinki. The volunteers signed an informed consent, completed a standard health questionnaire, and were screened for their habitual physical activity. None had any previous history of neuromuscular disorders. Volunteers included those involved in regular strength-training programs [strength-training group (ST), $n = 8$; age, 22.2 (2.5) yr [mean (SD)]; body mass, 85.2 (8.3) kg; height 181.2 (9.3) cm] and control individuals who were only involved in light to moderate aerobic activity [control group, $n = 8$; age, 23.4 (3.1) yr; body mass, 73.2 (7.5) kg; height 177.3 (7.5) cm]. All volunteers were students from the Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy. Before the first familiarization session, the volunteers were asked to report their physical activity habits. ST volunteers were required to be in a strength-training program that targeted strength and power for at least 3 yr and for a minimum of three times per week. The training programs were classical models of progressive strength training for enhancing muscle strength and power that targets all major upper and lower muscle groups. The training protocols performed by the ST cohort closely matched the guidelines reported by American College of Sports Medicine (2) for enhancing strength and power. The individuals from the ST cohort also performed national and international competitions

that involved explosive tasks, such as volleyball (4 subjects), javelin throw (1 subject), boxing (1 subject), and karate (1 subject). Controls were required to be involved in moderate to light aerobic exercise less than twice per week and were not involved in any form of regular strength or power training. All subjects were instructed to avoid strenuous exercise and caffeine for 48 and 24 h, respectively, before their visit to the laboratory.

Study Overview

Participants visited the laboratory on two occasions, 1 wk apart. During the first visit, they performed a familiarization test to become acquainted with the experimental protocol. The familiarization session included elbow flexion maximal voluntary isometric contractions (MVC) and maximal voluntary isometric explosive contractions of their dominant arm (self-reported). During the second visit, the volunteers performed the experimental session with concurrent recordings of force (MVC and isometric explosive force contractions) and high-density surface electromyography (HDsEMG).

Measurements

Force recording. Both the familiarization and measurement sessions were conducted using an isokinetic dynamometer (KIN-COM dynamometer; Chattecx, Chattanooga, TN) with the elbow of the dominant limb flexed to 90°. The reliability and feasibility of this dynamometer have been described previously (31) and used in previous studies that assessed maximal RTD (1). The chair configuration was established during the familiarization session and replicated in the main trial. Waist and shoulder straps were tightly fastened to prevent extraneous movements. The waist strap was fastened across the pelvis, and two other straps were fastened across the shoulders (39). This setup was comfortable and well tolerated by the participants of the study. The shoulder was in a neutral position with the upper arm parallel to the trunk (humerus in a pendent position), and the forearm was midway between pronation and supination. The elbow joint was secured in a padded brace with Velcro straps. The dynamometer load cell (KIN-COM dynamometer; Chattecx) consisted of four strain gauges. The wrist strap was consistently secured to the styloid process of the radius and was in series with a calibrated linear response from the dynamometer load cell that was positioned perpendicular to the radius. The center of the lever arm was aligned to the distal lateral epicondyle of the humerus. Subsequently, the lever arm length was measured as the distance between the distal lateral epicondyle of the humerus and the styloid process of the radius. The analog force signal was amplified and sampled at 2,048 Hz with an external analog-to-digital (A/D) converter (EMG-USB2+; OT Bioelettronica, Turin, Italy). Two personal computers recorded the data with the software OT BioLab (OT Bioelettronica) and LabVIEW 8.0 (National Instruments, Austin, TX). The force signal was displayed for visual feedback during the tests. Force signals were corrected for the effect of gravity.

HDsEMG recordings. Two bidimensional arrays (matrices) of 64 electrodes each [dimensions for 1 matrix: 1-mm diameter, 8-mm interelectrode distance, 13 rows (10.9 cm) $\times$ 5 columns (3.7 cm); gold-coated; OT Bioelettronica; Fig. 1A] were used for recording HDsEMG signals. The skin was treated by shaving, light abrasion, and cleansing with 70% ethanol. An experienced investigator identified the muscle belly of the biceps brachii (BB) through palpation, and a surgical marker was used to delineate the perimeter of the muscle. Before placing the electrodes, the arm circumference and the skinfold thickness (Harpender skinfold caliper) were measured. Successively, both matrices were placed over the BB using biadhesive foams (Spes Medica, Battipaglia, Italy). The grids were mounted close to each other to form an array of 128 electrodes (Fig. 1A). The array was located between the proximal and distal regions of the BB, along the direction of the muscle fibers, covering most of the BB area (30; Fig. 1A). The large number of electrodes used allowed the accurate




Fig. 1. *A*: two high-density surface electromyogram (EMG) arrays of 64 electrodes each. *B*: time-torque curve during an isometric explosive contraction (black curve) and the activity of 128 monopolar channels recorded from the biceps brachii muscle. *C*: 10 single-differential EMG signals during an explosive force contraction from a control subject. The innervation zone (IZ) and several motor unit action potentials (MUAPs) propagating in the distal direction can be seen. *D*: in these signals recorded from a strength-trained individual, several MUAPs propagating at a significant higher velocity can be seen compared with the control subject. The dashed curve indicates the time-torque curve. A.U., arbitrary units.

identification of the innervation zone and selection of channels with propagating action potentials. Moreover, the high-density configuration improves the reliability of MFCV estimates and of the EMG recordings considerably (30, 50). The ground electrode was placed on the wrist of the nondominant arm. Two reference electrodes were placed on the vertebra prominens and on the acromion. The EMG signals were amplified, band-pass filtered, and converted to digital data by a multichannel amplifier (3-dB bandwidth, 10–500 Hz; EMG-USB2+ multichannel amplifier; OT Bioelettronica). The same multichannel amplifier synchronized the HDsEMG and force signals.

Procedures

Before the measurements, the volunteers performed a standardized warm-up, which consisted of four contractions at 50% of their perceived MVF, four at ~75%, and one submaximal ~90% contraction. Each contraction was separated by 15 s. Following the warm-up with a recovery time of 5 min, the subjects completed three MVC, with 1 min of rest in between. The volunteers were encouraged to “push as hard as possible” for at least 3 s while receiving feedback on the exerted force and on the force exerted in the previous MVC. The greatest force was recorded as the MVF. After 5 min of rest, eight explosive force contractions were performed in two blocks of four contractions each. The blocks were separated by 5 min of rest, and the individual contractions within each block were separated by 20 s of rest. For the explosive force contractions, volunteers were instructed to relax and flex the elbow “as fast and hard” as possible after hearing an auditory cue. Volunteers were instructed to exceed the 75% of MVF threshold, which was displayed with a horizontal cursor on the monitor, without performing any countermovement. Only contractions that did not show any countermovement (≤ 0.5 N from the baseline of force in the 150 ms before the onset of force) were included within each block and used for the analysis.

Force signal analysis. In the off-line analysis, the analog force signal was converted to Newtons and multiplied by the respective lever arm to obtain torque (N·m). Successively, the torque signal was filtered with a low-pass fourth-order, zero-lag Butterworth filter with a cutoff frequency of 400 Hz. The torque onset (T_0) was determined with a visual detection method used in previous studies (51). After the onset detection was found, the torque was filtered with a zero-lag Butterworth filter with a cutoff frequency of 20 Hz. This two-step filtering procedure allowed us to first detect precisely the onset of torque in the 400-Hz low-pass-filtered signal (52), and then the 20-Hz filter removed all the nonphysiological frequencies in the signal. The 20-Hz low-pass filter guaranteed an undistorted signal in all cases with respect to the 400-Hz filtered signal, which was checked in all the explosive torque contractions. The onset value was used to determine

the torque values at 50 ms (T_{50}), 100 ms (T_{100}), 150 ms (T_{150}), and 200 ms (T_{200}) after the onset (Fig. 1, C and D). The five contractions with the highest torque values at T_{150} were used for the analysis. The absolute torque values at different time points were normalized by the respective maximal torque [$T/\text{maximal voluntary torque (MVT)}$]. The RTD (i.e., $\text{RTD}_{100-150} = T_{150} - T_{100}/0.05$ s) was calculated for three time intervals, RTD_{0-50} , RTD_{50-100} , and $\text{RTD}_{100-150}$, for both the absolute and relative torque values (e.g., $\text{RTD}_{\text{rel}_{0-50}}$; Fig. 2, A and B). The torque analysis was completed with MATLAB 2015 (The MathWorks, Natick, MA).

EMG processing. Single-differential HDsEMG signals were calculated from the monopolar derivations for each column of the two bidimensional arrays. Single-differential HDsEMG signals for each column were visually inspected, and the six single-differential HDsEMG signal channels with the highest coefficient of correlation (CC; $\text{CC} \geq 0.8$) and clear motor unit action potential propagation without shape change from the nearest innervation zone to the distal tendon (Fig. 1, C and D) were chosen for the analysis. The columns of the matrix that were selected for the estimates of MFCV corresponded to the central part (between the first 3 central columns) of the two bidimensional arrays, as they corresponded to the channels with the highest quality (CC and propagation). MFCV was computed using an algorithm that allows highly accurate estimates of conduction velocities from multichannel EMG signals and whose reliability and validity have been previously assessed in both isometric and dynamic contractions, with robust intraclass CC ($>75\%$ intraclass CC; 27, 29, 30). During isometric contractions the between-day coefficient of variability in MFCV is lower than 2%, with intraclass CC $>88\%$ (40). The use of ≥ 6 EMG channels allowed us to detect changes in MFCV as small as 0.1 m/s compared with estimates from a pair of bipolar signals (>0.4 m/s; 22). The BB muscle was chosen in this study because it has been shown to provide estimates of MFCV with good reliability during submaximal (50% MVC) steady-state contractions (30). It has been shown that MFCV can be accurately assessed also during dynamic explosive tasks (47). Maximal muscle fiber conduction velocity (MFCV_{MAX}) was estimated during the MVF contraction in time windows of 50 ms, from 500 ms before to 250 ms after peak torque during the MVF. The choice of this interval for the estimation of MFCV_{MAX} was due to the delay between recruitment and motor unit peak twitch forces. After determining the maximal value of MFCV, MFCV was estimated during the explosive torque contractions in intervals of 50 ms. For each explosive contraction, the onset of EMG activity was assessed visually (Fig. 1, C and D), and MFCV was estimated from five intervals, corresponding to the electromechanical delay [$\text{EMD} = T_0$ (s) – EMG onset (s)] and the four 50-ms intervals following T_0 (MFCV_{EMD} , MFCV_{0-50} , MFCV_{50-100} ,




Fig. 2. Rate of torque development in absolute (RTD, A) and normalized (RTD-rel, B) values. Black bars represent the strength-trained individuals, and white bars represent the controls. Data are reported as means (SD). * $P < 0.001$.

MFCV_{100–150}, and MFCV_{150–200}). MFCV absolute and normalized values [MFCV (m/s)/MFCV_{MAX}, e.g., MFCV-rel_{0–50}] were averaged over the five explosive contractions selected for the analysis.

Statistics

The Shapiro-Wilk test confirmed the normal distribution of the extracted variables. The number of participants needed for the study was estimated with a statistical power analysis test (function “sampsizepr” in MATLAB) using previous data on MFCV and RTD and successively progressively tested with the data collected in the present study. The significance level of the power test was set with a P value of 0.05. Two-way repeated-measures analysis of variance (group $\times$ time) was used to assess differences in explosive force, RTD and MFCV, for both absolute and normalized values. The analysis of variance included the five time intervals from EMG or torque onset during the explosive contractions. Bonferroni stepwise corrected paired t -tests were used to assess differences between groups at different time intervals. Moreover, Pearson product-moment CC was used to assess the linear relation between RTD and MFCV for each individual cohort, and the coefficient of determination (R^2) was used as an index of prediction power. The Bonferroni correction was applied to the regression significance values. Independent-sample t -tests were used to assess differences between groups for all other variables (MVF, MFCV_{MAX}, skinfold thickness, and arm circumference). Statistical analysis was completed using SPSS version 14 (SPSS, Chicago, IL) and MATLAB. The significance level was set at $P < 0.05$. Data are reported as means (SD).

RESULTS

EMD, Anthropometry, and Statistical Power

The EMD did not differ between the two groups [ST 62.42 (10.81) vs. controls 60.24 (12.12) ms, $P > 0.05$]. Moreover, there was no difference in the subcutaneous fat layer thickness between groups [ST 4.11 (0.71), control group 4.45 (1.05) mm, $P > 0.05$] as assessed by skinfold measures, whereas the arm circumference was greater for ST [36.01 (1.51) vs. 29.3 (2.58) cm, $P < 0.05$]. The power analysis indicated that seven subjects per cohort were needed to obtain a power of 90% for MFCV and RTD estimates.

Torque

Maximal torque was significantly greater for ST compared with controls [99.64 (21.62) vs. 60.56 (8.74) N·m, $P < 0.001$].

ST also developed higher torques at 50, 100, 150, and 200 ms from contraction onset ($P < 0.001$). When torque at different time points was expressed relative to the MVT, ST achieved higher relative torques in the first two phases of contraction [T_{50} and T_{100} ; T_{50} , 43.84 (5.29) vs. 17.12 (4.61)%MVT; T_{100} , 63.45 (10.02) vs. 33.25 (4.42)%MVT, $P < 0.001$]. The relative explosive torque at T_{150} and T_{200} was similar for both cohorts ($P > 0.05$).


The absolute RTD during the initial phase (T_{50}) of the contraction was greater for the resistance-trained individuals than for controls [861.82 (104.60) vs. 342.05 (93.08) N·m/s, $P < 0.001$; Fig. 2A]. The absolute RTD over the consecutive time windows did not differ (Fig. 2A). In addition, the normalized RTD was greater only in the first 50 ms of contraction for ST [887.66 (152.08) vs. 568.54 (148.66)%MVT/s, $P < 0.001$; Fig. 2B]. Conversely, the normalized RTD_{50–100} and RTD_{100–150} was greater for the controls [536.48 (112.08) vs. 404.91 (96.36)%MVT/s, 356.21 (110.48) vs. 271.41 (69.44)%MVT/s, $P < 0.001$; Fig. 2B].

Muscle Fiber Conduction Velocity

Maximal MFCV at MVT (MFCV_{MAX}) ranged from 5.05 to 5.82 m/s in ST and from 4.93 to 5.26 m/s in the controls with the ST group having a significantly higher MFCV_{MAX} compared with controls [5.37 (0.27) vs. 5.04 (0.11) m/s, $P < 0.001$].

MFCV during the explosive torque contractions (from the EMD to T_{200}) ranged from 3.44 to 5.45 m/s (and the lowest value of MFCV corresponded to the time interval during the EMD, MFCV_{EMD}) when grouping all participants. The average MFCV values across subjects at each time window can be observed for the two cohorts in Fig. 3. The MFCV value consistently increased in the second time interval (MFCV_{0–50}; Fig. 3, $P < 0.01$). This indicates recruitment of motor units with progressively larger diameter fibers with increase in torque (10, 11, 41), which is related to the progressive recruitment by size (11, 37). However, the ST achieved a higher MFCV_{EMD} compared with the controls [4.44 (0.13) vs. 3.83 (0.20) m/s, $P < 0.001$; Fig. 3A], even when MFCV_{EMD} was normalized to MFCV_{MAX} [82.57 (3.13) vs. 75.86 (3.55) MFCV-rel_{EMD}, $P < 0.001$; Fig. 3B]. Moreover, the early phase of absolute and

Fig. 3. Muscle fiber conduction velocity in absolute (MFCV, A) and normalized (MFCV-rel, B) values. ●, Strength-trained individuals. Data are reported as means (SD). EMD, electromechanical delay; EMG, electromyogram; MAX, maximum. * $P < 0.01$.



normalized MFCV estimates was correlated to RTD (Fig. 4; see RESULTS, *RTD and MFCV Correlations*).

The ST cohort maintained greater absolute and normalized MFCV values throughout the full duration of the explosive contractions ($P < 0.001$; Fig. 3A). Interestingly, the time-MFCV curve had a similar pattern for ST compared with the controls (Fig. 3, A and B). MFCV indeed increased linearly from EMD until reaching a plateau at MFCV_{50-100} ($P < 0.001$) for both groups (Fig. 3, A and B). This observation indirectly indicates that the muscle full motor unit recruitment may have been completed before the first 100 ms of explosive torque production.

RTD and MFCV Correlations

The estimates of MFCV were positively correlated with RTD only in the time window RTD_{0-50} (Fig. 4). For the resistance-trained individuals, a negative correlation was found between normalized $\text{RTD}_{100-150}$ and absolute MFCV [average R^2 for all MFCV estimates when plotted as a function of $\text{RTD}_{100-150}$ values = -0.59 (0.24), $P < 0.001$]. During the same RTD time window, the correlation for the controls was not significant ($P > 0.05$).

DISCUSSION

MFCV was measured during explosive force contractions in a group of resistance-trained individuals and a control cohort. ST exhibited greater explosive torque, early RTD, and greater MFCV with respect to controls throughout the contraction. When explosive torque was normalized to maximal torque, ST had a higher RTD at the beginning of the contraction (0–50 ms). Moreover, a greater absolute and normalized conduction velocity (MFCV_{MAX}) during the EMD and in the first 50 ms of torque generation was observed for the ST group. This result indicates a recruitment of motor units with greater conduction velocities. This is the first study showing that ST may recruit larger motor units in a shorter amount of time.

Muscle Fiber Conduction Velocity

The average MFCV values are in agreement with previous reports of MFCV during steady-state contractions. For example, Farina et al. reported estimates of MFCV in the BB during isometric steady-state contractions at 50% MVC of ~ 4.6 m/s (30). Zwarts and Arendt-Nielsen estimated MFCV at high contraction forces of the BB and reported values ranging between 3.22 and 5.11 m/s (57). MFCV average values in the present study were also in agreement with estimates of single-

motor unit conduction velocities using intramuscular EMG recordings during voluntary and electrical activation of the BB muscle. Moreover, the present estimates are also in accordance with other studies involving different muscular contractions and protocols (25, 26, 32, 41, 42, 47, 48).

Interestingly, only two studies assessed MFCV in power athletes and only during electrical stimulation and maximal voluntary contractions (43, 49). Sadoyama et al. reported a significantly higher maximal MFCV in a group of trained sprinters compared with endurance runners (4.84 vs. 4.31 m/s; 49). Moreover, they reported a significant relation between the relative area of fast-twitch fibers and conduction velocity (49).

Recently, Methenitis et al. estimated MFCV during electrical stimulation of muscle fibers in endurance runners and power-trained and ST individuals and separately measured RTD (43). They reported significant relations between MFCV, muscle fiber cross-sectional area, and rate of force development (43). However, estimates of MFCV were assessed during electrical stimulation and thus separately from the voluntary generation of explosive force. Therefore, it was not possible to associate the underlying neural strategies of muscle control with explosive force performance. Collectively, these previous results indicate that MFCV may be an indicator of muscle explosive performance, although no previous study assessed MFCV during explosive torque generation.

Explosive Torque and RTD

The RTD was significantly greater for ST during the early phase of explosive torque generation (Fig. 3A). However, when the moment-time curve was normalized to the maximal strength, the RTD for the resistance-trained subjects was significantly different only in the first 50 ms of the contraction (Fig. 3B). Because the relative explosive torque at 150 and 200 ms from contraction onset was similar between the two groups, the controls developed higher RTD during the second and third time windows from force onset (Fig. 3B).

Previous studies found an increase in the EMG amplitude and rate of force development in the initial phase of contraction after 4 wk of explosive training (7). In addition, a greater normalized rate of force development in the first 50 ms of contraction was found for power athletes during knee extensor explosive torque (51). Because the first 50 ms of contraction strongly reflect neural activation (13, 15, 16, 54), strength or power training presumably increases RTD by a faster recruitment of motor units, as discussed below.




Fig. 4. Correlation between the rate of torque development during the first 50 ms of contraction (RTD_{0-50}) and muscle fiber conduction velocity during the electromechanical delay (MFCV_{EMD} , A) and in the first 50 ms of contraction (MFCV_{0-50} , B). ●, Control individuals. Correlation coefficients (R^2) and regression lines are given. * $P < 0.05$.

MFCV During the Explosive Phase of Contraction

ST individuals have the ability to develop higher levels of force in the first 50 ms of contraction. This seems to be associated with greater MFCV in the same time interval, which indicates recruitment of motor units with greater conduction velocity. The role of motor unit recruitment during explosive force contraction is not well understood because it is not possible to identify representative populations of motor units in very short time intervals.

The primary determinant of motor unit twitch force is the number of muscle fibers innervated by the axon (17, 53). Motor unit peak twitch forces in humans range from ~6 to ~78 mN·m with maximal tetanic forces ranging from ~30 to ~200 mN·m (36, 38). Therefore, one of the mechanisms that determined the increase in RTD during the first 0–50-ms interval in the ST may have been the recruitment of larger motor units with greater and faster twitches. There is evidence showing correlations between electrically evoked twitch torque and early voluntary rate of force development, which could be associated with the differences in muscle fiber composition and/or Ca^{2+} saturation for the trained individuals (3, 34).

Interestingly, the two groups showed similar EMD values, which is in accordance with a previous study that compared the EMD in power-trained and untrained cohorts (51). This finding is, however, contradictory with a higher MFCV during the EMD. A higher absolute MFCV value should theoretically anticipate the release of Ca^{2+} and thus the rise in force. However, these problems may be related to the techniques employed in assessing the delay between the neural and muscular apparatus. The EMD during explosive contractions may not be sensitive to differences in neural activation due to a compressed recruitment (12, 15). We have recently shown that when the EMD is assessed as the time difference between the neural drive and force during the sustained contractions, the central nervous system modulates the delay broadly and according to the rate of force development (12). Indeed, the neuromechanical delay seems to be predominantly influenced by the type of recruited motor units and the intrinsic properties of the motor neuron (5, 12). Future studies assessing the neuromechanical delay in strength/power-trained individuals may be warranted.

MFCV increases with voluntary force production due to the relation between motor unit recruitment thresholds and fiber diameter (4, 8, 11, 35). This association implies that the ordered recruitment of motor units may be assessed by estimates of conduction velocity (10). We have recently reported that large, high-threshold motor units innervate fibers with large diameter (11), which explains the association between motor unit mechanical properties and conduction velocity previously reported (4). Moreover, we have recently demonstrated that the increase in average MFCV during voluntary force contractions is associated with the progressive recruitment of motor units with increasing conduction velocity and predicts recruitment thresholds at the individual subject level (10).

In the present study, MFCV was the average of the conduction velocities of the active motor units during explosive force contractions, in time intervals of 50 ms following EMG onset. We showed that there may be significant differences in the recruited motor units during explosive tasks in ST compared with moderately active individuals. Absolute MFCV values

were greater in ST throughout the full duration of the explosive contractions (Fig. 3A). Moreover, the early absolute and normalized MFCV were positively associated with RTD (Fig. 4). Because absolute MFCV values are linearly related to the diameters of muscle fibers, higher absolute conduction velocity values may indicate that ST have muscle fibers with larger diameters due to the strength training-induced hypertrophy (43) compared with controls (24, 43).

However, when MFCV values were normalized to the maximal value during MVF (full motor unit recruitment), ST had a significantly greater MFCV-rel during the initial phase of the explosive contractions. Specifically, MFCV-rel_{EMD} and MFCV-rel_{0–50} were on average ~9% greater (Fig. 4B). This suggests that during the early phase of explosive force, ST have the ability to recruit motor units with faster conduction velocities in a shorter time. It takes ~100 ms more for the controls to reach similar values of normalized MFCV compared with the ST group. Interestingly, the changes in conduction velocity did not differ between groups (Fig. 4, A and B), and the MFCV plateaued in the interval 50–100 ms, which can be interpreted as full motor unit recruitment (10). This interpretation is in agreement with previous studies reporting that most motor units are recruited at one-third of maximal force during explosive contractions (15, 16). Moreover, MFCV increased in all subjects from the EMD to 0–50 ms, indicating that the ordered recruitment according to the size of the motor unit was preserved during the explosive tasks in both groups.

The underlying mechanism that may determine an increase in explosive force for the ST individuals may be an anticipated recruitment of high-threshold motor units with high conduction velocities. The difference between MFCV of high- and low-threshold motor units within a muscle is ~2 m/s (11). A faster motor unit recruitment (and conduction velocity) would achieve greater peak mechanical torques in a shorter time. The release of calcium from the sarcoplasmic reticulum is correlated to the speed of the action potential on the fiber membrane (21). Indeed, MFCV is related to motor unit time-to-peak twitch forces (4). The increase in MFCV may potentially allow a faster calcium uptake and thus anticipating the rise in force.

Van Cutsem et al. reported an increase in motor unit discharge rates following ballistic training (55) and concluded, in agreement with other studies, that RTD depends on motor unit discharge rate (16, 18, 55). On the other hand, the recruitment threshold of motor units significantly influences the discharge rate at a given absolute force (18). Anticipating the recruitment of high-threshold motor units would result in reaching motor unit peak discharge rate and motor unit peak RTD in a shorter time. Accordingly, in the present study, MFCV was positively associated with RTD (Fig. 4A), suggesting that motor unit recruitment may play an important role in explosive force production.

Interestingly, the correlation between RTD_{0–50} and early MFCV values (MFCV_{EMD,0–50}) was different for the ST and untrained individuals. MFCV_{0–50} was not correlated with RTD_{0–50} in the ST group (Fig. 4). This result indicates that ST completed the motor unit recruitment during the very early phase of explosive force, i.e., between the EMD and the first 50 ms from contraction onset (Fig. 4B). The increase in MFCV during the explosive force at the 50- and 100-ms time points for the ST is presumably due to some subjects continuing the

recruitment while the subjects with higher RTD achieved a faster plateau in MFCV (Fig. 4). Indeed, it took more time for the untrained individuals to reach high MFCV (and full motor unit recruitment) values (Figs. 3 and 4).

It must be noted that the number of subjects in the present study may be too low for a correlation study. Moreover, the cross-sectional design cannot isolate the innate and environmental factors that contributed to the explosive force and MFCV differences found between the cohorts. However, in the present study the trained individuals performed combined strength and explosive training for >3 yr. Recent evidence showed a significant increase in explosive force production after 12 wk of isometric explosive training compared with isometric sustained-contraction strength training (6). The large differences in the early RTD for the trained subjects in the present study may also indicate that the neural contributors to explosive strength could be related to chronic exposure to explosive/strength training and that the neural adaptations may continue over time. Future studies assessing the neural contributors to explosive force in large cohorts and longitudinal (>1 yr) interventions are warranted.

Conclusion

Resistance-trained individuals showed higher RTD and explosive force in the very early phase of contraction that was accompanied by an increase in absolute and normalized MFCV compared with controls. These observations may be explained by recruitment of fast-twitch motor units (i.e., large motor units with large muscle fiber diameters) in a shorter amount of time in the resistance-trained cohort than in controls. In addition to the functional implications in the study of human explosive force, the study also presents a methodology that may be applied in the assessment of the neural strategies of muscle control in health, training, and pathology.

GRANTS

A. Del Vecchio has received funding from the University of Rome "Foro Italico." F. Negro has received funding from the European Union's Horizon 2020 Programme for Research and Innovation under Marie Skłodowska-Curie Grant Agreement 702491 ["Connectivity in the neural control of muscles in stroke patients" (NeuralCon)].

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

AUTHOR CONTRIBUTIONS

A.D.V., F.N., D. Falla, I.B., D. Farina, and F.F. conceived and designed research; A.D.V. performed experiments; A.D.V. analyzed data; A.D.V., F.N., D. Falla, I.B., and D. Farina interpreted results of experiments; A.D.V. prepared figures; A.D.V. drafted manuscript; A.D.V., F.N., D. Falla, I.B., D. Farina, and F.F. edited and revised manuscript; A.D.V., F.N., D. Falla, I.B., D. Farina, and F.F. approved final version of manuscript.

REFERENCES

1. Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *J Appl Physiol* (1985) 93: 1318–1326, 2002. doi:10.1152/jappphysiol.00283.2002.
2. American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 41: 687–708, 2009. doi:10.1249/MSS.0b013e3181915670.
3. Andersen LL, Aagaard P. Influence of maximal muscle strength and intrinsic muscle contractile properties on contractile rate of force development. *Eur J Appl Physiol* 96: 46–52, 2006. doi:10.1007/s00421-005-0070-z.
4. Andreassen S, Arendt-Nielsen L. Muscle fibre conduction velocity in motor units of the human anterior tibial muscle: a new size principle parameter. *J Physiol* 391: 561–571, 1987. doi:10.1113/jphysiol.1987.sp016756.
5. Baldissera F, Cavallari P, Cerri G. Motoneuronal pre-compensation for the low-pass filter characteristics of muscle. A quantitative appraisal in cat muscle units. *J Physiol* 511: 611–627, 1998. doi:10.1111/j.1469-7793.1998.611bh.x.
6. Balshaw TG, Massey GJ, Maden-Wilkinson TM, Tillin NA, Folland JP. Training-specific functional, neural, and hypertrophic adaptations to explosive- vs. sustained-contraction strength training. *J Appl Physiol* (1985) 120: 1364–1373, 2016. doi:10.1152/jappphysiol.00091.2016.
7. Barry BK, Warman GE, Carson RG. Age-related differences in rapid muscle activation after rate of force development training of the elbow flexors. *Exp Brain Res* 162: 122–132, 2005. doi:10.1007/s00221-004-2127-3.
8. Blijham PJ, ter Laak HJ, Schelhaas HJ, van Engelen BGM, Stegeman DF, Zwarts MJ. Relation between muscle fiber conduction velocity and fiber size in neuromuscular disorders. *J Appl Physiol* (1985) 100: 1837–1841, 2006. doi:10.1152/jappphysiol.01009.2005.
9. Del Vecchio A, Bazzucchi I, Felici F. Variability of estimates of muscle fiber conduction velocity and surface EMG amplitude across subjects and processing intervals. *J Electromyogr Kinesiol* 40: 102–109, 2018. doi:10.1016/j.jelekin.2018.04.010.
10. Del Vecchio A, Negro F, Felici F, Farina D. Associations between motor unit action potential parameters and surface EMG features. *J Appl Physiol* (1985) 123: 835–843, 2017. doi:10.1152/jappphysiol.00482.2017.
11. Del Vecchio A, Negro F, Felici F, Farina D. Distribution of muscle fiber conduction velocity for representative samples of motor units in the full recruitment range of the tibialis anterior muscle. *Acta Physiol (Oxf)* 38: e12930, 2018. doi:10.1111/apha.12930.
12. Del Vecchio A, Ubada A, Sartori M, Azorin JM, Felici F, Farina D. The central nervous system modulates the neuromechanical delay in a broad range for the control of muscle force. *J Appl Physiol* (1985). In press. doi:10.1152/jappphysiol.00135.2018.
13. de Ruiter CJ, Kooistra RD, Paalman MI, de Haan A. Initial phase of maximal voluntary and electrically stimulated knee extension torque development at different knee angles. *J Appl Physiol* (1985) 97: 1693–1701, 2004. doi:10.1152/jappphysiol.00230.2004.
14. de Ruiter CJ, Vermeulen G, Toussaint HM, de Haan A. Isometric knee-extensor torque development and jump height in volleyball players. *Med Sci Sports Exerc* 39: 1336–1346, 2007. doi:10.1097/mss.0b013e318063c719.
15. Desmedt JE, Godaux E. Ballistic contractions in man: characteristic recruitment pattern of single motor units of the tibialis anterior muscle. *J Physiol* 264: 673–693, 1977. doi:10.1113/jphysiol.1977.sp011689.
16. Desmedt JE, Godaux E. Ballistic contractions in fast or slow human muscles: discharge patterns of single motor units. *J Physiol* 285: 185–196, 1978. doi:10.1113/jphysiol.1978.sp012566.
17. Dideriksen JL, Farina D. Motor unit recruitment by size does not provide functional advantages for motor performance. *J Physiol* 591: 6139–6156, 2013. doi:10.1113/jphysiol.2013.262477.
18. Duchateau J, Baudry S. Maximal discharge rate of motor units determines the maximal rate of force development during ballistic contractions in human. *Front Hum Neurosci* 8: 234, 2014. doi:10.3389/fnhum.2014.00234.
19. Duchateau J, Semmler JG, Enoka RM. Training adaptations in the behavior of human motor units. *J Appl Physiol* (1985) 101: 1766–1775, 2006. doi:10.1152/jappphysiol.00543.2006.
20. Erskine RM, Fletcher G, Folland JP. The contribution of muscle hypertrophy to strength changes following resistance training. *Eur J Appl Physiol* 114: 1239–1249, 2014. doi:10.1007/s00421-014-2855-4.
21. Farina D, Arendt-Nielsen L, Graven-Nielsen T. Effect of temperature on spike-triggered average torque and electrophysiological properties of low-threshold motor units. *J Appl Physiol* (1985) 99: 197–203, 2005. doi:10.1152/jappphysiol.00059.2005.
22. Farina D, Arendt-Nielsen L, Merletti R, Graven-Nielsen T. Assessment of single motor unit conduction velocity during sustained contractions of the tibialis anterior muscle with advanced spike triggered averaging. *J Neurosci Methods* 115: 1–12, 2002. doi:10.1016/S0165-0270(01)00510-6.

23. Farina D, Cescon C, Negro F, Enoka RM. Amplitude cancellation of motor-unit action potentials in the surface electromyogram can be estimated with spike-triggered averaging. *J Neurophysiol* 100: 431–440, 2008. doi:10.1152/jn.90365.2008.
24. Farina D, Ferguson RA, Macaluso A, De Vito G. Correlation of average muscle fiber conduction velocity measured during cycling exercise with myosin heavy chain composition, lactate threshold, and $\dot{V}O_{2\max}$. *J Electromyogr Kinesiol* 17: 393–400, 2007. doi:10.1016/j.jelekin.2006.03.003.
25. Farina D, Gazzoni M, Camelia F. Conduction velocity of low-threshold motor units during ischemic contractions performed with surface EMG feedback. *J Appl Physiol* (1985) 98: 1487–1494, 2005. doi:10.1152/jappphysiol.01032.2004.
26. Farina D, Macaluso A, Ferguson RA, De Vito G. Effect of power, pedal rate, and force on average muscle fiber conduction velocity during cycling. *J Appl Physiol* (1985) 97: 2035–2041, 2004. doi:10.1152/jappphysiol.00606.2004.
27. Farina D, Muhammad W, Fortunato E, Meste O, Merletti R, Rix H. Estimation of single motor unit conduction velocity from surface electromyogram signals detected with linear electrode arrays. *Med Biol Eng Comput* 39: 225–236, 2001. doi:10.1007/BF02344807.
28. Farina D, Negro F, Dideriksen JL. The effective neural drive to muscles is the common synaptic input to motor neurons. *J Physiol* 592: 3427–3441, 2014. doi:10.1113/jphysiol.2014.273581.
29. Farina D, Pozzo M, Merlo E, Bottin A, Merletti R. Assessment of average muscle fiber conduction velocity from surface EMG signals during fatiguing dynamic contractions. *IEEE Trans Biomed Eng* 51: 1383–1393, 2004. doi:10.1109/TBME.2004.827556.
30. Farina D, Zagari D, Gazzoni M, Merletti R. Reproducibility of muscle-fiber conduction velocity estimates using multichannel surface EMG techniques. *Muscle Nerve* 29: 282–291, 2004. doi:10.1002/mus.10547.
31. Farrell M, Richards JG. Analysis of the reliability and validity of the kinetic communicator exercise device. *Med Sci Sports Exerc* 18: 44–49, 1986. doi:10.1249/00005768-198602000-00009.
32. Felici F, Bazzucchi I, Sgrò P, Quinzi F, Conti A, Aversa A, Gizzi L, Mezzullo M, Romanelli F, Pasquali R, Lenzi A, Di Luigi L. Acute severe male hypo-testosteronemia affects central motor command in humans. *J Electromyogr Kinesiol* 28: 184–192, 2016. doi:10.1016/j.jelekin.2015.12.004.
33. Felici F, Rosponi A, Sbriccoli P, Filligoi GC, Fattorini L, Marchetti M. Linear and non-linear analysis of surface electromyograms in weightlifters. *Eur J Appl Physiol* 84: 337–342, 2001. doi:10.1007/s004210000364.
34. Folland JP, Buckthorpe MW, Hannah R. Human capacity for explosive force production: neural and contractile determinants. *Scand J Med Sci Sports* 24: 894–906, 2014. doi:10.1111/sms.12131.
35. Håkansson CH. Conduction velocity and amplitude of the action potential as related to circumference in the isolated fibre of frog muscle. *Acta Physiol Scand* 37: 14–34, 1956. doi:10.1111/j.1748-1716.1956.tb01338.x.
36. Heckman CJ, Enoka RM. Motor unit. *Compr Physiol* 2: 2629–2682, 2012. doi:10.1002/cphy.c100087.
37. Henneman E. Relation between size of neurons and their susceptibility to discharge. *Science* 126: 1345–1347, 1957. doi:10.1126/science.126.3287.1345.
38. Macefield VG, Fuglevand AJ, Bigland-Ritchie B. Contractile properties of single motor units in human toe extensors assessed by intraneural motor axon stimulation. *J Neurophysiol* 75: 2509–2519, 1996. doi:10.1152/jn.1996.75.6.2509.
39. Maffiuletti NA, Aagaard P, Blazevich AJ, Folland J, Tillin N, Duchateau J. Rate of force development: physiological and methodological considerations. *Eur J Appl Physiol* 116: 1091–1116, 2016. doi:10.1007/s00421-016-3346-6.
40. Martinez-Valdes E, Laine CM, Falla D, Mayer F, Farina D. High-density surface electromyography provides reliable estimates of motor unit behavior. *Clin Neurophysiol* 127: 2534–2541, 2016. doi:10.1016/j.clinph.2015.10.065.
41. Masuda T, De Luca CJ. Recruitment threshold and muscle fiber conduction velocity of single motor units. *J Electromyogr Kinesiol* 1: 116–123, 1991. doi:10.1016/1050-6411(91)90005-P.
42. Merletti R, Knaflitz M, De Luca CJ. Myoelectric manifestations of fatigue in voluntary and electrically elicited contractions. *J Appl Physiol* (1985) 69: 1810–1820, 1990. doi:10.1152/jappl.1990.69.5.1810.
43. Methenitis S, Karandreas N, Spengos K, Zaras N, Stasinaki AN, Terzis G. Muscle fiber conduction velocity, muscle fiber composition, and power performance. *Med Sci Sports Exerc* 48: 1761–1771, 2016. doi:10.1249/MSS.0000000000000954.
44. Nandedkar SD, Stålberg E. Simulation of single muscle fibre action potentials. *Med Biol Eng Comput* 21: 158–165, 1983. doi:10.1007/BF02441531.
45. Peñaillillo L, Blazevich A, Numazawa H, Nosaka K. Rate of force development as a measure of muscle damage. *Scand J Med Sci Sports* 25: 417–427, 2015. doi:10.1111/sms.12241.
46. Plonsey R, Barr RC. *Bioelectricity: A Quantitative Approach*. New York: Springer, 2007.
47. Pozzo M, Merlo E, Farina D, Antonutto G, Merletti R, Di Prampero PE. Muscle-fiber conduction velocity estimated from surface EMG signals during explosive dynamic contractions. *Muscle Nerve* 29: 823–833, 2004. doi:10.1002/mus.20049.
48. Rodriguez-Falces J, Malanda A, Latasa I, Lavilla-Oiz A, Navallas J. Influence of timing variability between motor unit potentials on M-wave characteristics. *J Electromyogr Kinesiol* 30: 249–262, 2016. doi:10.1016/j.jelekin.2016.08.003.
49. Sadoyama T, Masuda T, Miyata H, Katsuta S. Fibre conduction velocity and fibre composition in human vastus lateralis. *Eur J Appl Physiol Occup Physiol* 57: 767–771, 1988. doi:10.1007/BF01076001.
50. Staudenmann D, Kingma I, Stegeman DF, van Dieën JH. Towards optimal multi-channel EMG electrode configurations in muscle force estimation: a high density EMG study. *J Electromyogr Kinesiol* 15: 1–11, 2005. doi:10.1016/j.jelekin.2004.06.008.
51. Tillin NA, Jimenez-Reyes P, Pain MT, Folland JP. Neuromuscular performance of explosive power athletes versus untrained individuals. *Med Sci Sports Exerc* 42: 781–790, 2010. doi:10.1249/MSS.0b013e3181be9c7e.
52. Tillin NA, Pain MT, Folland JP. Identification of contraction onset during explosive contractions. Response to Thompson et al. “Consistency of rapid muscle force characteristics: influence of muscle contraction onset detection methodology” [*J Electromyogr Kinesiol* 2012; 22(6):893–900]. *J Electromyogr Kinesiol* 23: 991–994, 2013. doi:10.1016/j.jelekin.2013.04.015.
53. Tötösy de Zepetnek JE, Zung HV, Erdebil S, Gordon T. Innervation ratio is an important determinant of force in normal and reinnervated rat tibialis anterior muscles. *J Neurophysiol* 67: 1385–1403, 1992. doi:10.1152/jn.1992.67.5.1385.
54. Van Cutsem M, Duchateau J. Preceding muscle activity influences motor unit discharge and rate of torque development during ballistic contractions in humans. *J Physiol* 562: 635–644, 2005. doi:10.1113/jphysiol.2004.074567.
55. Van Cutsem M, Duchateau J, Hainaut K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. *J Physiol* 513: 295–305, 1998. doi:10.1111/j.1469-7793.1998.295by.x.
56. Vila-Chã C, Falla D, Farina D. Motor unit behavior during submaximal contractions following six weeks of either endurance or strength training. *J Appl Physiol* (1985) 109: 1455–1466, 2010. doi:10.1152/jappphysiol.01213.2009.
57. Zwarts MJ, Arendt-Nielsen L. The influence of force and circulation on average muscle fibre conduction velocity during local muscle fatigue. *Eur J Appl Physiol Occup Physiol* 58: 278–283, 1988. doi:10.1007/BF00417263.